

THE SYMMETRIC FUNCTION

$$h^0(\overline{M}_{0,n}, \mathcal{L}_1^{x_1} \otimes \mathcal{L}_2^{x_2} \otimes \cdots \otimes \mathcal{L}_n^{x_n})$$

RAHUL PANDHARIPANDE¹

0. INTRODUCTION

0.1. Summary of Results. Let $n \geq 3$ be an integer. Let $(C, p_1, \dots, p_n)$ be a reduced, connected, (at worst) nodal curve of arithmetic genus 0 with n nonsingular marked points. $(C, p_1, \dots, p_n)$ is *Deligne-Mumford stable* if $w_C(p_1 + \dots + p_n)$ is ample (where w_C is the dualizing sheaf). Let $\overline{M}_{0,n}$ denote the fine moduli space of Deligne-Mumford stable, n -pointed, genus 0 curves. A foundational treatment of $\overline{M}_{0,n}$ can be found in [Kn]. Let $S_n = \{1, 2, \dots, n\}$ be the marking set. For each $i \in S_n$, a line bundle $\mathcal{L}_i$ on $\overline{M}_{0,n}$ is obtained by the following prescription: the fiber of $\mathcal{L}_i$ at the moduli point $[C, p_1, \dots, p_n] \in \overline{M}_{0,n}$ is the cotangent space T_{C, p_i}^* .

Proposition 1. *Let $x_1, x_2, \dots, x_n$ be non-negative integers.*

$$\forall k > 0, \quad H^k(\overline{M}_{0,n}, \mathcal{L}_1^{x_1} \otimes \mathcal{L}_2^{x_2} \otimes \cdots \otimes \mathcal{L}_n^{x_n}) = 0.$$

For non-negative x_i , let $\gamma_n(x_1, x_2, \dots, x_n) = h^0(\overline{M}_{0,n}, \mathcal{L}_1^{x_1} \otimes \mathcal{L}_2^{x_2} \otimes \cdots \otimes \mathcal{L}_n^{x_n})$. By Proposition (1) and the Hirzebruch-Riemann-Roch Theorem, γ_n is a polynomial function: $\gamma_n \in \mathbb{Q}[x_1, x_2, \dots, x_n]$. The symmetric group Σ_n acts naturally on $\overline{M}_{0,n}$ by permuting the markings. The action of Σ_n permutes the isomorphism classes of the line bundles $\mathcal{L}_i$ in the obvious manner. Therefore, γ_n is a *symmetric* polynomial in $x_1, x_2, \dots, x_n$. Since $\overline{M}_{0,n}$ is $n - 3$ dimensional, γ_n is of degree (at most) $n - 3$.

Date: 15 November 1995.

¹Research partially supported by an NSF Post-Doctoral Fellowship.

Let $\sigma_1, \sigma_2, \dots, \sigma_n$ denote the elementary symmetric functions in $x_1, x_2, \dots, x_n$.

$$\begin{aligned}\sigma_1 &= x_1 + \dots + x_n \\ \sigma_2 &= x_1x_2 + x_1x_3 + \dots + x_{n-1}x_n \\ &\vdots \\ \sigma_n &= x_1x_2 \cdots x_n.\end{aligned}$$

The following equivalence is well-known:

$$\mathbb{Q}[\sigma_1, \dots, \sigma_n] = \left(\mathbb{Q}[x_1, \dots, x_n] \right)^{\Sigma_n}.$$

Hence, $\gamma_n \in \mathbb{Q}[\sigma_1, \dots, \sigma_n]$.

Let $R = \mathbb{Q}[\sigma_1, \sigma_2, \dots]$ be the infinite polynomial ring in the variables $\{\sigma_d\}_{d \in \mathbb{N}^+}$. Define a grading on R by assigning to the variable σ_d the weight d . For $f \in R$, the *degree* of f is the highest weight of the monomials that appear in f .

A $\mathbb{Q}$ -linear transformation $T : R \rightarrow R$ is defined as follows. Let $f \in R$. Let $e = \text{degree}(f)$. Therefore, $f \in \mathbb{Q}[\sigma_1, \sigma_2, \dots, \sigma_e]$. The element f corresponds to a symmetric function in the variables $x_1, \dots, x_m$ provided $m \geq e$. Let $g(x_1, \dots, x_m)$ be defined for non-negative integers x_i by

$$g(x_1, \dots, x_m) = f(x_1, \dots, x_m) + \sum_{i=1}^m \sum_{j=0}^{x_i-1} f(x_1, \dots, x_{i-1}, j, x_{i+1}, \dots, x_m). \quad (1)$$

The function g is manifestly a symmetric polynomial in $x_1, \dots, x_m$ of degree (at most) $e + 1$. Therefore $g \in \mathbb{Q}[\sigma_1, \sigma_2, \dots, \sigma_e, \sigma_{e+1}]$. The element g is independent of m provided $m \geq e+1$. Define $T(f) = g \in R$ for m in the stable range $m \geq e + 1$.

Proposition 2. *The function $\gamma_n \in \mathbb{Q}[\sigma_1, \dots, \sigma_n]$ is determined by:*

$$\gamma_n = T^{n-3}(1).$$

The author has benefited from conversations with W. Fulton, M. Kapranov, and M. Thaddeus.

0.2. Computing T . Computing T is tedious but straightforward. If $f = 1$, then $\text{degree}(f) = 0$. $T(1)$ can be computed in the variable x_1 .

By equation (1),

$$g(x_1) = f(x_1) + \sum_{j=0}^{x_1-1} f(j) = 1 + x_1 = 1 + \sigma_1.$$

Hence $T(1) = 1 + \sigma_1$.

Let $n = 4$. $\overline{M}_{0,4} \cong \mathbb{P}^{\mathbb{A}^k}$. Via this isomorphism, $\mathcal{L}_i \cong \mathcal{O}_{\mathbb{P}^{\mathbb{A}^k}}(1)$ for $1 \leq i \leq 4$. The function γ_4 is easily evaluated:

$$\begin{aligned} \gamma_4(x_1, x_2, x_3, x_4) &= h^0(\mathbb{P}^{\mathbb{A}^k}, \mathcal{O}_{\mathbb{P}^{\mathbb{A}^k}}(\curvearrowright_{\mathbb{A}^k} + \curvearrowright_{\mathbb{A}^k} + \curvearrowright_{\mathbb{A}^k} + \curvearrowright_{\mathbb{A}^k})) \\ &= 1 + x_1 + x_2 + x_3 + x_4 \\ &= 1 + \sigma_1 \\ &= T(1). \end{aligned}$$

Next, let $f = \sigma_1$. $T(\sigma_1)$ can be computed in the variables x_1, x_2 .

$$\begin{aligned} g(x_1, x_2) &= f(x_1, x_2) + \sum_{j=0}^{x_1-1} f(j, x_2) + \sum_{j=0}^{x_2-1} f(x_1, j) \\ &= x_1 + x_2 + \frac{1}{2}(x_1^2 - x_1) + x_1x_2 + \frac{1}{2}(x_2^2 - x_2) + x_1x_2 \\ &= \frac{1}{2}\sigma_1 + \frac{1}{2}\sigma_1^2 + \sigma_2 \end{aligned}$$

Therefore, $T(\sigma_1) = \frac{1}{2}\sigma_1 + \frac{1}{2}\sigma_1^2 + \sigma_2$. By Proposition (2),

$$\gamma_5 = T^2(1) = T(\sigma_1 + 1) = T(\sigma_1) + T(1) = 1 + \frac{3}{2}\sigma_1 + \frac{1}{2}\sigma_1^2 + \sigma_2.$$

Proposition (2) leads to an easy calculation of $h^0(\overline{M}_{0,n}, \mathcal{L}_1^{x_1})$. By relation (1), for $n \geq 4$,

$$h^0(\overline{M}_{0,n}, \mathcal{L}_1^{x_1}) = \sum_{j=0}^{x_1} h^0(\overline{M}_{0,n-1}, \mathcal{L}_1^{x_1}).$$

It is easily seen $h^0(\overline{M}_{0,n}, \mathcal{L}_1^{x_1}) = \binom{n-3+x_1}{x_1}$ uniquely satisfies the above recursion and the boundary conditions at $n = 3$. The global sections of $\mathcal{L}_1^{x_1}$ can also be computed by examining the morphism $\overline{M}_{0,n} \rightarrow \mathbb{P}^{\mathbb{A}^k - \mathbb{A}^k}$ determined by $H^0(\overline{M}_{0,n}, \mathcal{L}_1)$ (see section (1.1)).

The function γ_n is tabulated below for small values of n :

$$\gamma_3 = 1$$

$$\gamma_4 = 1 + \sigma_1$$

$$\gamma_5 = 1 + \frac{3}{2}\sigma_1 + \frac{1}{2}\sigma_1^2 + \sigma_2$$

$$\gamma_6 = 1 + \frac{11}{6}\sigma_1 + \sigma_1^2 + \sigma_2 + \frac{1}{6}\sigma_1^3 + \sigma_1\sigma_2 + 2\sigma_3$$

$$\begin{aligned} \gamma_7 = 1 + \frac{25}{12}\sigma_1 + \frac{35}{24}\sigma_1^2 + \frac{5}{4}\sigma_2 + \frac{5}{12}\sigma_1^3 + \frac{5}{4}\sigma_1\sigma_2 - \frac{1}{4}\sigma_3 \\ + \frac{1}{24}\sigma_1^4 + \frac{1}{2}\sigma_1^2\sigma_2 + \frac{5}{2}\sigma_1\sigma_3 + \frac{1}{4}\sigma_2^2 + \frac{11}{2}\sigma_4 \end{aligned}$$

$$\begin{aligned} \gamma_8 = 1 + \frac{137}{60}\sigma_1 + \frac{15}{8}\sigma_1^2 + \frac{5}{4}\sigma_2 + \frac{17}{24}\sigma_1^3 + \frac{11}{6}\sigma_1\sigma_2 + \frac{11}{4}\sigma_3 \\ + \frac{1}{8}\sigma_1^4 + \frac{3}{4}\sigma_1^2\sigma_2 - \frac{3}{4}\sigma_1\sigma_3 + \frac{1}{4}\sigma_2^2 - \frac{21}{2}\sigma_4 \\ + \frac{1}{120}\sigma_1^5 + \frac{1}{6}\sigma_1^3\sigma_2 + \frac{3}{2}\sigma_1^2\sigma_3 + \frac{1}{4}\sigma_1\sigma_2^2 + \frac{17}{2}\sigma_1\sigma_4 + \sigma_2\sigma_3 + 19\sigma_5 \end{aligned}$$

1. THE LINE BUNDLES $\mathcal{L}_i$

1.1. Expressions for $\mathcal{L}_i$. Let $\rho : U_n \rightarrow \overline{M}_{0,n}$ be the universal family of pointed, stable curves. There are n sections $s_1, \dots, s_n$ of ρ corresponding to the markings. Consider the marking i . The projection ρ takes s_i isomorphically to $\overline{M}_{0,n}$. Let

$$x = [C, p_1, \dots, p_n] \in \overline{M}_{0,n}.$$

Let $y \in s_i$ satisfy $\rho(y) = x$. The normal bundle to s_i in U_n at y is canonically the tangent space T_{C,p_i} . Therefore

$$c_1(\mathcal{L}_i) \cong \rho_*(-s_i^2). \tag{2}$$

Note U_n is canonically isomorphic to $\overline{M}_{0,n+1}$. $\mathcal{L}_{n+1}$ on U_n can be expressed as follows:

$$\mathcal{L}_{n+1} \cong \omega_\rho(s_1 + \dots + s_n) \quad (3)$$

where ω_ρ is the relative dualizing sheaf of the morphism $\rho : U_n \cong \overline{M}_{0,n+1} \rightarrow \overline{M}_{0,n}$. The proof of (3) requires a square diagram:

$$\begin{array}{ccc} \overline{M}_{0,n+1\#} & \xrightarrow{\pi_{n+1}} & \overline{M}_{0,n+1} \\ \nu \downarrow & & \rho \downarrow \\ \overline{M}_{0,n\#} & \xrightarrow{\pi_n} & \overline{M}_{0,n} \end{array}$$

$\overline{M}_{0,n+1\#}$ is the moduli space of $n+2$ -pointed genus 0 stable curves with marking set $\{1, \dots, n+1, \#\}$. The morphisms ν , ρ , π_n , and π_{n+1} are all contraction maps. Let $D_{n+1,\#}$ be the boundary divisor of $\overline{M}_{0,n+1\#}$ corresponding to the partition

$$\{n+1, \#\} \cup \{1, \dots, n\}.$$

$D_{n+1,\#}$ is the section of π_{n+1} corresponding to the marking $n+1$. A simple examination of the blow-ups involved in the above square yields:

$$\pi_{n+1}^*(\omega_\rho(s_1 + \dots + s_n))|_{D_{n+1,\#}} \cong \omega_\nu|_{D_{n+1,\#}}.$$

Since $\omega_\nu|_{D_{n+1,\#}} \cong -D_{n+1,\#}|_{D_{n+1,\#}}$, (3) is established by (2).

Another method of viewing $\mathcal{L}$ is as follows. Fix n marked points in general linear position in $\mathbb{P}^{\times-\#}$. M. Kapranov has shown the closure in the Hilbert scheme of the locus of rational normal curves passing through the n marked points is canonically $\overline{M}_{0,n}$ ([K]). The universal curve over the Hilbert scheme restricts to U_n over $\overline{M}_{0,n}$. Since the universal curve over the Hilbert scheme naturally maps to $\mathbb{P}^{\times-\#}$, a morphism

$$\mu : U_n \rightarrow \mathbb{P}^{\times-\#}$$

is obtained. U_n is canonically isomorphic to $\overline{M}_{0,n+1}$. There is an isomorphism:

$$\mathcal{L}_{n+1} \cong \mu^*(\mathcal{O}_{\mathbb{P}^{\times-\#}}(1)). \quad (4)$$

On each fiber of $\rho : U_n \rightarrow \overline{M}_{0,n}$, $\mu^*(\mathcal{O}_{\mathbb{P}^{\times-\#}}(1))$ is isomorphic to $\omega_\rho(s_1 + \dots + s_n)$ (see [K]). Since both line bundles $\mu^*(\mathcal{O}_{\mathbb{P}^{\times-\#}}(1))$ and $\omega_\rho(s_1 + \dots + s_n)$ are trivial on the sections s_i of ρ ,

$$\mu^*(\mathcal{O}_{\mathbb{P}^{\times-\#}}(1)) \cong \omega_\rho(s_1 + \dots + s_n)$$

on U_n . Now (4) follows from (3).

Finally, it is useful to express $\mathcal{L}_{n+1}$ on $\overline{M}_{0,n+1}$ as a linear combination of boundary divisors. Let $j, k \in S_n$ be distinct markings.

$$\mathcal{L}_{n+1} \cong \sum_{n+1 \in A \subset S_{n+1}, j, k \notin A} D_A \quad (5)$$

where the set A is a subset of the marking set S_{n+1} satisfying $|A| \geq 2$. As above, D_A is the boundary divisor corresponding to the partition $S_{n+1} = A \cup A^c$. Recall the map

$$\mu : \overline{M}_{0,n+1} \cong U_n \rightarrow \mathbb{P}^{\times -\#}.$$

Consider the unique hyperplane $H \subset \mathbb{P}^{\times -\#}$ passing through the $n-2$ marked points $S_n \setminus \{j, k\}$. The divisor $\mu^*(H)$ is easily seen to be the right side of (5). By (4), the isomorphism of (5) is established. See [W] for another proof of (5).

1.2. Contraction. It will be helpful in the sequel to denote $\mathcal{L}_i$ on $\overline{M}_{0,n}$ by $\mathcal{L}_{i,n}$. Let $n \geq 3$. Consider the contraction morphism

$$\rho : \overline{M}_{0,n+1} \rightarrow \overline{M}_{0,n}$$

obtained by omitting the marking $n+1$.

Lemma 1. $\rho^*(\mathcal{L}_{i,n}) + D_{i,n+1} \cong \mathcal{L}_{i,n+1}$.

Proof. Let $j, k \in S_n$ satisfy $j, k \neq i$. By the results of section (1.1)

$$\rho^*(\mathcal{L}_{i,n}) \cong \sum_{i \in A \subset S_n, j, k \notin A} \rho^*(D_A).$$

A comparison with

$$\mathcal{L}_{i,n+1} \cong \sum_{i \in A \subset S_{n+1}, j, k \notin A} D_A$$

yields Lemma (1). □

2. THE PROOF OF PROPOSITION (1)

2.1. The Induction Ladder. Consider the following (infinite) commutative diagram:

$$\begin{array}{ccccc} \overline{M}_{0,n+1\#} & \xrightarrow{\nu_n} & \overline{M}_{0,n\#} & \xrightarrow{\nu_{n-1}} & \overline{M}_{0,n-1\#} \\ \pi_{n+1} \downarrow & & \pi_n \downarrow & & \pi_{n-1} \downarrow \\ \overline{M}_{0,n+1} & \xrightarrow{\rho_n} & \overline{M}_{0,n} & \xrightarrow{\rho_{n-1}} & \overline{M}_{0,n-1} \end{array} \quad (6)$$

The maps are all contraction morphisms: π_n contracts $\#$, ν_{n-1} contracts n , ρ_{n-1} contracts n . The diagram starts with $\overline{M}_{0,3}$ on the lower right corner and extends left. Denote the composition $\rho_n \circ \cdots \circ \rho_{m-1}$ by $\rho_{m,n}$ (similarly for ν).

Recall $\mathcal{L}_{i,n}$ is the line bundle corresponding to the i^{th} marking on $\overline{M}_{0,n}$. For $m > n$, the pull-back line bundle on $\overline{M}_{0,m}$, $\rho_{m,n}^*(\mathcal{L}_{i,n})$, is also denoted by the same symbol $\mathcal{L}_{i,n}$.

Let $\mathcal{L}_{\#,n\#}$ be the line bundle corresponding to the marking $\#$ on $\overline{M}_{0,n\#}$. Again $\mathcal{L}_{\#,n\#}$ will also denote the pull-back line bundle, $\nu_{m,n}^*(\mathcal{L}_{\#,n\#})$ on $\overline{M}_{0,m\#}$ for $m > n$.

Proposition (1) is established by an induction on the ladder (6). The following Lemmas are needed in the induction.

Lemma 2. *Let $n \geq 3$. The line bundle $\mathcal{L}_{\#,n\#}$ behaves well under ladder base changes:*

- (i.) *For all $k \geq 0$ and $m \geq n$, $\pi_{m*}(\mathcal{L}_{\#,n\#}^k) \cong \rho_{m,n}^* \pi_{n*}(\mathcal{L}_{\#,n\#}^k)$.*
- (ii.) *For all $k \geq 0$ and $m \geq n$, $R^1 \pi_{m*}(\mathcal{L}_{\#,n\#}^k) = 0$.*

Proof. By (4), $\mathcal{L}_{\#,n\#}$ is generated by global sections on $\overline{M}_{0,n\#}$. Therefore $\mathcal{L}_{\#,n\#}^k$ is generated globally on $\overline{M}_{0,m\#}$. Any line bundle generated by global sections on a genus 0, stable, pointed curve has no higher cohomology. Hence, $\mathcal{L}_{\#,n\#}^k$ has no higher cohomology on the fibers of π_m . By the Cohomology and Base Change Theorems,

$$R^1 \pi_{m*}(\mathcal{L}_{\#,n\#}^k) = 0.$$

Consider the fiber product:

$$\tau_m : \overline{M}_{0,m} \times_{\overline{M}_{0,n}} \overline{M}_{0,n\#} \rightarrow \overline{M}_{0,m}.$$

Again by Base Change, $\tau_{m*}(\mathcal{L}_{\#,n\#}^k) \cong \rho_{m,n}^* \pi_{n*}(\mathcal{L}_{\#,n\#}^k)$. There is natural map

$$\overline{M}_{0,m\#} \rightarrow \overline{M}_{0,m} \times_{\overline{M}_{0,n}} \overline{M}_{0,n\#} \tag{7}$$

which commutes with π_m and τ_m . It is easy to check that the natural map of vector bundles $\tau_{m*}(\mathcal{L}_{\#,n\#}^k) \rightarrow \pi_{m*}(\mathcal{L}_{\#,n\#}^k)$ induced by (7) is an isomorphism on fibers. $\square$

Lemma 3. *Let $n \geq 4$. The following exact sequences of vector bundles exist on $\overline{M}_{0,n}$ for each pair of integers $a, b \geq 0$:*

$$0 \rightarrow \pi_{n*}(\mathcal{L}_{\#,n-1\#}^{a+1} \otimes \mathcal{L}_{\#,n\#}^b) \rightarrow \pi_{n*}(\mathcal{L}_{\#,n-1\#}^a \otimes \mathcal{L}_{\#,n\#}^{b+1}) \rightarrow \mathcal{L}_{n,n}^a \rightarrow 0. \quad (8)$$

Proof. By Lemma (1), there is a linear equivalence $\mathcal{L}_{\#,n-1\#} + D_{\#,n} \cong \mathcal{L}_{\#,n\#}$. Note:

$$\begin{aligned} \mathcal{L}_{\#,n\#}|_{D_{\#,n}} &= 0, \\ \mathcal{L}_{\#,n-1\#}|_{D_{\#,n}} &\cong -D_{\#,n}|_{D_{\#,n}}. \end{aligned} \quad (9)$$

Tensoring the sequence (10) with $\mathcal{L}_{\#,n-1\#}^a \otimes \mathcal{L}_{\#,n\#}^{b+1}$ yields (11).

$$0 \rightarrow \mathcal{O}(-D_{\#,n}) \rightarrow \mathcal{O} \rightarrow \mathcal{O}_{D_{\#,n}} \rightarrow 0 \quad (10)$$

$$0 \rightarrow \mathcal{L}_{\#,n-1\#}^{a+1} \otimes \mathcal{L}_{\#,n\#}^b \rightarrow \mathcal{L}_{\#,n-1\#}^a \otimes \mathcal{L}_{\#,n\#}^{b+1} \rightarrow \mathcal{L}_{\#,n-1\#}^a|_{D_{\#,n}} \rightarrow 0. \quad (11)$$

Equivalence (9) implies $\pi_{n*}(\mathcal{L}_{\#,n-1\#}^a|_{D_{\#,n}}) \cong \mathcal{L}_{n,n}^a$. By a global sections argument as in Lemma (2), the terms of sequence (11) have vanishing higher direct images under π_n . Sequence (8) is obtained by pushing-forward sequence (11). $\square$

Lemma 4. *Let $n \geq 3$. Let $\mathcal{N}$ be a line bundle on $\overline{M}_{0,n}$ satisfying the following condition for all non-negative integers $z_3, \dots, z_n$ and all $k > 0$:*

$$H^k(\overline{M}_{0,n}, \mathcal{N} \otimes \bigotimes_{i=3}^n \mathcal{L}_{i,i}^{z_i}) = 0. \quad (12)$$

Then for all $b \geq 0$ and $k > 0$,

$$H^k(\overline{M}_{0,n}, \pi_{n*}(\mathcal{L}_{\#,n\#}^b) \otimes \mathcal{N}) = 0. \quad (13)$$

Proof. If $b = 0$, the vanishing (13) is a consequence of (12) since $\pi_{n*}(\mathcal{O}) \cong \mathcal{O}$. Assume $b > 0$. The proof is a simple consequence of Lemmas (2) and (3). Consider the sequences for $0 \leq j \leq b-1$ obtained from (8) by tensoring with $\mathcal{N}$:

$$0 \rightarrow \pi_{n*}(\mathcal{L}_{\#,n-1\#}^{j+1} \otimes \mathcal{L}_{\#,n\#}^{b-j-1}) \otimes \mathcal{N} \rightarrow \pi_{n*}(\mathcal{L}_{\#,n-1\#}^j \otimes \mathcal{L}_{\#,n\#}^{b-j}) \otimes \mathcal{N} \rightarrow \mathcal{L}_{n,n}^j \otimes \mathcal{N} \rightarrow 0.$$

The vanishing (13) is reduced by these sequences and repeated application of (12) to:

$$\forall b \geq 0, \forall k > 0, \quad H^k(\overline{M}_{0,n}, \pi_{n*}(\mathcal{L}_{\#,n-1\#}^b) \otimes \mathcal{N}) = 0.$$

But since, $\pi_{n*}(\mathcal{L}_{\#,n-1\#}^b) \cong \rho_{n-1}^* \pi_{n-1*}(\mathcal{L}_{\#,n-1\#}^b)$ by Lemma (2), it suffices to show:

$$\forall b \geq 0, \forall k > 0, \quad H^k(\overline{M}_{0,n}, \rho_{n-1}^* \pi_{n-1*}(\mathcal{L}_{\#,n-1\#}^b) \otimes \mathcal{N}) = 0.$$

Pulling-back the sequences (8) for $n-1$ to $\overline{M}_{0,n}$ via ρ_{n-1} , tensoring with $\mathcal{N}$, and repeatedly applying (12), the vanishing (13) is further reduced to:

$$\forall b \geq 0, \forall k > 0, \quad H^k(\overline{M}_{0,n}, \rho_{n-1}^* \pi_{n-1*}(\mathcal{L}_{\#,n-2\#}^b) \otimes \mathcal{N}) = 0.$$

Applying Lemma (2), it suffices to show

$$\forall b \geq 0, \forall k > 0, \quad H^k(\overline{M}_{0,n}, \rho_{n,n-2}^* \pi_{n-2*}(\mathcal{L}_{\#,n-2\#}^b) \otimes \mathcal{N}) = 0.$$

This process finally reduces the original claim (13) to:

$$\forall b \geq 0, \forall k > 0, \quad H^k(\overline{M}_{0,n}, \rho_{n,3}^* \pi_{3*}(\mathcal{L}_{\#,3\#}^b) \otimes \mathcal{N}) = 0.$$

Since $\pi_{3*}(\mathcal{L}_{\#,3\#}^b)$ is a trivial bundle over the point $\overline{M}_{0,3}$, the Lemma is proven. $\square$

2.2. The Induction. A slightly stronger version of Proposition (1) is needed for the induction:

Proposition 1'. Let $n \geq 3$. Let $x_{i,j}$ (for $1 \leq i \leq j$, $3 \leq j \leq n$) be non-negative integers.

$$\forall k > 0, \quad H^k(\overline{M}_{0,n}, \bigotimes_{i,j} \mathcal{L}_{i,j}^{x_{i,j}}) = 0.$$

Proof. The Proposition certainly holds for $n=3$ since $\dim(\overline{M}_{0,3}) = 0$. Assume the Proposition holds for n . Let $x_{i,j}$ (for $1 \leq i \leq j$, $3 \leq j \leq n$) be non-negative integers. Let $y_1, \dots, y_n, y_{\#}$ also be non-negative integers. Let $\mathcal{L}_{1,n\#}, \dots, \mathcal{L}_{n,n\#}, \mathcal{L}_{\#,n\#}$ denote the line bundles on $\overline{M}_{0,n\#}$ corresponding to the markings $1, \dots, n, \#$. Consider the following line bundle on $\overline{M}_{0,n\#}$:

$$\mathcal{L}(y_1, \dots, y_n, y_{\#}) \otimes \mathcal{N} \cong \mathcal{L}_{\#,n\#}^{y_{\#}} \otimes \bigotimes_{t=1}^n \mathcal{L}_{t,n\#}^{y_t} \otimes \bigotimes_{i,j} \mathcal{L}_{i,j}^{x_{i,j}}.$$

where $\mathcal{N} = \bigotimes_{i,j} \mathcal{L}_{i,j}^{x_{i,j}}$ and

$$\mathcal{L}(y_1, \dots, y_n, y_{\#}) = \mathcal{L}_{\#,n\#}^{y_{\#}} \otimes \bigotimes_{t=1}^n \mathcal{L}_{t,n\#}^{y_t}.$$

It suffices to prove for all $k > 0$, $H^k(\overline{M}_{0,n\#}, \mathcal{L}(y_1, \dots, y_n, y_{\#}) \otimes \mathcal{N}) = 0$.

As before, let π_n denote the contraction map $\pi_n : \overline{M}_{0,n\#} \rightarrow \overline{M}_{0,n}$. For each $1 \leq t \leq n$,

$$\mathcal{L}_{t,n} + D_{t,\#} \cong \mathcal{L}_{t,n\#}.$$

The following exact sequences on $\overline{M}_{0,n\#}$ therefore exist for each $1 \leq t \leq n$ and non-negative integers a, b :

$$0 \rightarrow \mathcal{L}_{t,n}^{a+1} \otimes \mathcal{L}_{t,n\#}^b \rightarrow \mathcal{L}_{t,n}^a \otimes \mathcal{L}_{t,n\#}^{b+1} \rightarrow \mathcal{L}_{t,n}^a|_{D_{t,\#}} \rightarrow 0. \quad (14)$$

Via the natural identification $\pi_n : D_{t,\#} \cong \overline{M}_{0,n}$, the following isomorphisms hold:

$$\begin{aligned} \mathcal{L}_{\#,n\#}|_{D_{t,\#}} &\cong \mathcal{O}, \\ \mathcal{L}_{t,n\#}|_{D_{t,\#}} &\cong \mathcal{O}, \\ \forall s \neq t, \mathcal{L}_{s,n\#}|_{D_{t,\#}} &\cong \mathcal{L}_{s,n}, \\ \forall 1 \leq i \leq j, 3 \leq j \leq n, \mathcal{L}_{i,j}|_{D_{t,\#}} &\cong \mathcal{L}_{i,j}. \end{aligned} \quad (15)$$

By repeated use of (14) along with the inductive assumption of cohomology vanishing on $\overline{M}_{0,n} \cong D_{t,\#}$, it suffices to prove

$$\forall k > 0, \quad H^k(\overline{M}_{0,n\#}, \mathcal{L}_{\#,n\#}^{y\#} \otimes \mathcal{N}') = 0,$$

$$\mathcal{N}' \cong \bigotimes_{i=1}^n \mathcal{L}_{i,n}^{y_i} \otimes \mathcal{N}.$$

In effect, the sequences (14) are used to convert factors of $\mathcal{L}_{i,n\#}$ to $\mathcal{L}_{i,n}$ one at a time. The remaining factors of $\mathcal{L}_{\#,n\#}$ are handled by the following method. Since $R^1\pi_{n*}(\mathcal{L}_{\#,n\#}^{y\#} \otimes \mathcal{N}') = 0$, for all $k > 0$,

$$H^k(\overline{M}_{0,n\#}, \mathcal{L}_{\#,n\#}^{y\#} \otimes \mathcal{N}') \cong H^k(\overline{M}_{0,n}, \pi_{n*}(\mathcal{L}_{\#,n\#}^{y\#} \otimes \mathcal{N}')).$$

Since $\pi_{n*}(\mathcal{L}_{\#,n\#}^{y\#} \otimes \mathcal{N}') \cong \pi_{n*}(\mathcal{L}_{\#,n\#}^{y\#}) \otimes \mathcal{N}'$ and $\mathcal{N}'$ satisfies the vanishing (12) by the inductive assumption on $\overline{M}_{0,n}$, the proof of the Proposition is completed by Lemma (4). $\square$

3. THE PROOF OF PROPOSITION (2)

The proof of Proposition (1') gives a recursive (in n) method to calculate

$$h^0(\overline{M}_{0,n}, \bigotimes_{i,j} \mathcal{L}_{i,j}^{x_{i,j}})$$

for non-negative $x_{i,j}$. There are 3 reasons why this full recursion is not pursued here:

- (i.) The full recursion is complicated.

- (ii.) More data is required in the recursion than is needed in Proposition (2).
- (iii.) The full recursion does not respect the symmetry of the variables in Proposition (2).

Fortunately there is simple way to sidestep these problems.

Recall $\gamma_n(x_1, x_2, \dots, x_n) = h^0(\overline{M}_{0,n}, \mathcal{L}_{1,n}^{x_1} \otimes \mathcal{L}_{2,n}^{x_2} \otimes \dots \otimes \mathcal{L}_{n,n}^{x_n})$ is a symmetric function of degree (at most) $n - 3$ in $x_1, \dots, x_n$. The map from symmetric functions in $x_1, \dots, x_n$ to symmetric functions in $x_1, \dots, x_{n-1}$ given by setting $x_n = 0$ is bijective for symmetric functions of degree at most $n - 1$. Both spaces are spanned by monomials in $\sigma_1, \dots, \sigma_{n-1}$ of degree at most $n - 1$. Hence γ_n is completely determined by $\gamma_n(x_1, \dots, x_{n-1}, 0)$.

Suppose the symmetric function γ_n is known. To determine γ_{n+1} , it suffices to know $\gamma_{n+1}(x_1, \dots, x_n, 0)$. The vanishing of the last entry leads to a great simplification in the recursion of Proposition (1'). Let

$$\gamma_{n+1}(x_1, \dots, x_n, 0) = h^0(\overline{M}_{0,n\#}, \bigotimes_{t=1}^n \mathcal{L}_{t,n\#}^{x_t}).$$

From the sequences (for $1 \leq t \leq n$)

$$0 \rightarrow \mathcal{L}_{t,n}^{a+1} \otimes \mathcal{L}_{t,n\#}^b \rightarrow \mathcal{L}_{t,n}^a \otimes \mathcal{L}_{t,n\#}^{b+1} \rightarrow \mathcal{L}_{t,n}^a|_{D_{t,\#}} \rightarrow 0 \quad (16)$$

and the restriction equations (15), the following relation is easily deduced:

$$\gamma_{n+1}(x_1, \dots, x_n, 0) = \gamma_n(x_1, \dots, x_n) + \sum_{i=1}^n \sum_{j=0}^{x_i-1} \gamma_n(x_1, \dots, x_{i-1}, j, x_{i+1}, \dots, x_n).$$

The sequences (16), as before, are used to convert factors of $\mathcal{L}_{t,n\#}$ to $\mathcal{L}_{t,n}$. By Proposition (1'), all higher cohomology vanishes. It is the omission of the factor $\mathcal{L}_{\#,n\#}$ that simplifies the recursion. This concludes the proof of Proposition (2).

REFERENCES

- [K] M. Kapranov, *Veronese Curves And Grothendieck-Knudsen Moduli Spaces* $\overline{M}_{0,n}$, J. Alg. Geom. Vol. **2** No. **2** (1993), 239-262.
- [Kn] F. Knudsen, *Projectivity of the Moduli Space of Stable Curves II*, Math Scand. **52** (1983), 1225-1265.

- [W] E. Witten, *Two Dimensional Quantum Gravity And Intersection Theory On Moduli Space*, Surveys In Diff. Geom. **1** (1991), 243-310.

Department of Mathematics
University of Chicago
5743 S. University Ave
Chicago, IL 60637
rahul@math.uchicago.edu